arXiv:math/9807028v1 [math.QA] 7 Jul 1998

A class of non-symmetric solutions for the integrability condition of the Knizhnik-Zamolodchikov equation: a Hopf algebra approach

G. Militaru
University of Bucharest
Faculty of Mathematics
Str. Academiei 14
RO-70109 Bucharest 1, Romania
e-mail: gmilit@al.math.unibuc.ro

Abstract

Let M be a vector space over a field k and $R \in \mathrm{End}_k(M \otimes M)$. This paper studies what shall be called the Long equation: that is, the system of nonlinear equations $R^{12}R^{13} = R^{13}R^{12}$ and $R^{12}R^{23} = R^{23}R^{12}$ in $\mathrm{End}_k(M \otimes M \otimes M)$. Any symmetric solution of this system supplies us a solution of the integrability condition of the Knizhnik-Zamolodchikov equation: $[R^{12}, R^{13}+R^{23}] = 0$ ([4] or [10]). We shall approach this equation by introducing a new class of bialgebras, which we call Long bialgebras: these are pairs (H, σ), where H is a bialgebra and $\sigma : H \otimes H \to k$ is a k-bilinear map satisfying certain properties. The main theorem of this paper is a FRT type theorem: if M is finite dimensional, any solution R of the Long equation has the form $R = R_\sigma$, where M has a structure of a right comodule over a Long bialgebra $(L(R), \sigma)$, and R_σ is the special map $R_\sigma(m \otimes n) = \sum \sigma(m_{<1>} \otimes n_{<1>}) m_{<0>} \otimes n_{<0>}$.

0 Introduction

Let M be a finite dimensional vector space over a field k and $R \in \mathrm{End}_k(M \otimes M)$. In this paper we shall describe all solutions of what we have called the Long equation, namely the system of nonlinear equations

$$\begin{cases} R^{12}R^{13} = R^{13}R^{12} \\ R^{12}R^{23} = R^{23}R^{12} \end{cases} \tag{1}$$

in $\mathrm{End}_k(M \otimes M \otimes M)$. Our approach is similar to the one used by Faddeev, Reshetikhin and Takhtadjian in relating the solutions of the quantum Yang-Baxter equation to comodules over co-quasitriangular bialgebras (see [2]).

We shall introduce a new class of bialgebras which we shall call Long bialgebras. They are pairs (H, σ), where H is a bialgebra and $\sigma : H \otimes H \to k$ is a linear map satisfying the conditions $(L1) - (L5)$ from definition 3.3. The conditions $(L2) - (L5)$ are identical with the conditions $(B2) - (B5)$ from the definition of co-quasitriangular bialgebras. What differentiates Long bialgebras from the co-quasitriangular bialgebras is the condition $(L1)$ versus $(B1)$. This new class of bialgebras will play a fundamental role in solving the Long equation. More precisely, if (M, ρ) is a right comodule over a Long bialgebra (H, σ), then the special map

$$R_\sigma : M \otimes M \to M \otimes M, \quad R_\sigma(m \otimes n) = \sum \sigma(m_{<1>} \otimes n_{<1>}) m_{<0>} \otimes n_{<0>}$$

is a solution of the Long equation. Conversely, the main theorem of this paper is a FRT type theorem: if M is a finite dimensional vector space and R is a solution of the Long equation, then there exists a Long bialgebra $(L(R), \sigma)$ such that M has a structure of right $L(R)$-comodule and $R = R_\sigma$. Let us now look at the Long equation from a different angle, having in mind Radford's version of the FRT theorem ([9]): in the finite dimensional case, any solution R of the quantum Yang-Baxter equation has the form $R = R_{(M, \cdot, \rho)}$, where $(M, \cdot, \rho) \in {}_{A(R)}\mathcal{YD}^{A(R)}$, the category of Yetter-Drinfel'd modules. This is obtained immediately, keeping in mind the fact that a comodule (M, ρ) over a co-quasitriangular bialgebra (like $(A(R), \sigma)$) has a structure of Yetter-Drinfel'd module via

$$h \cdot m = \sum \sigma(m_{<1>} \otimes h) m_{<0>}$$

for all $h \in A(R)$, $m \in M$ and $R_{(M, \cdot, \rho)} = R_\sigma$. A similar phenomenon happens to the comodules over a Long bialgebra (H, σ): they become objects in the category ${}_H\mathcal{L}^H$ of H-dimodules (see Proposition 3.6). This category has been introduced by Long ([5]) for the case of a commutative and cocommutative H and studied in connection with the Brauer group of an H-dimodule algebra. For this reason, we called system (1) the Long equation and the new class of bialgebras, Long bialgebras.

Finally, we shall point out a connection between the symmetric solutions of the Long equation (e.g. the map R^ϕ from proposition 2.4) and solutions of the integrability condition of the Knizhnik-Zamolodchikov equation. Suppose that R is a solution of the Long equation; then R satisfies the equation $[R^{12}, R^{13} + R^{23}] = 0$, which is called the integrability condition of the Knizhnik-Zamolodchikov equation (see [10]). Moreover, we assume that R is symmetric, that is $R^{12} = R^{21}$. W is a solution of the Knizhnik-Zamolodchikov equation if it is a solution of the following system of differential equations:

$$\frac{\partial W}{\partial z^i} = h \sum_{i \neq j} \left(\frac{R^{ij}}{z^i - z^j}\right) W. \tag{2}$$

This is the equation for a function $W(z)$ taking values in $M^{\otimes n}$, representing a covariant constant section of the trivial bundle $Y_n \times M^{\otimes n} \to Y_n$ with flat connection $\sum_{i \neq j} R^{ij} \frac{dz^i}{z^i - z^j}$ (here k is the complex field $\mathbf{C}$, h is a complex parameter and $Y_n = \mathbf{C}^n$/multidiagonal). In other words, R is describing a family of flat connections on bundles with fiber $M^{\otimes n}$. For further details we refer to [4] and [10].

1 Preliminaries

Throughout this paper, k will be a field. All vector spaces, algebras, coalgebras and bialgebras considered are over k. $\otimes$ and Hom will mean $\otimes_k$ and Hom_k. For a coalgebra C, we will use Sweedler's Σ-notation, that is, $\Delta(c) = \sum c_{(1)} \otimes c_{(2)}$, $(I \otimes \Delta)\Delta(c) = \sum c_{(1)} \otimes c_{(2)} \otimes c_{(3)}$, etc. We will also use Sweedler's notation for right C-comodules: $\rho_M(m) = \sum m_{<0>} \otimes m_{<1>}$, for any $m \in M$ if (M, ρ_M) is a right C-comodule. $\mathcal{M}^C$ will be the category of right C-comodules and C-colinear maps and ${}_A\mathcal{M}$ will be the category of left A-modules and A-linear maps, if A is a k-algebra. An important role in the present paper will be played by $\mathcal{M}^n(k)$, the comatrix coalgebra of order n, i.e. $\mathcal{M}^n(k)$ is the n^2-dimensional vector space with $\{c_{ij} \mid i, j = 1, \cdots, n\}$ a k-basis such that

$$\Delta(c_{jk}) = \sum_{u=1}^{n} c_{ju} \otimes c_{uk}, \quad \varepsilon(c_{jk}) = \delta_{jk} \tag{3}$$

for all $j, k = 1, \cdots, n$. We view $T(\mathcal{M}^n(k))$ with the unique bialgebra structure which can be defined on the tensor algebra $T(\mathcal{M}^n(k))$, which extends the comultiplication Δ and the counity ε of $\mathcal{M}^n(k)$.

Let H be a bialgebra. An H-*dimodule* over H is a triple $(M, \cdot, \rho)$, where $(M, \cdot)$ is a left H-module, (M, ρ) is a right H-comodule such that the following compatibility condition holds:

$$\rho(h \cdot m) = \sum h \cdot m_{<0>} \otimes m_{<1>} \tag{4}$$

for all $h \in H$ and $m \in M$. The category of H-dimodules over H and H-linear H-colinear maps will be denoted by ${}_H\mathcal{L}^H$. This category was introduced for a commutative and cocommutative H by Long in [5].

For a vector space M, $\tau : M \otimes M \to M \otimes M$ will denote the flip map, that is $\tau(m \otimes n) = n \otimes m$ for all m, $n \in M$. If $R : M \otimes M \to M \otimes M$ is a linear map, we denote by R^{12}, R^{13}, R^{23} the maps of $\mathrm{End}_k(M \otimes M \otimes M)$ given by

$$R^{12} = R \otimes I, \quad R^{23} = I \otimes R, \quad R^{13} = (I \otimes \tau)(R \otimes I)(I \otimes \tau).$$

2 The Long equation

We shall start with the following:

Definition 2.1 *Let M be a vector space and $R \in \mathrm{End}_k(M \otimes M)$. We shall say that R is a solution for the Long equation if*

$$\begin{cases} R^{12}R^{13} = R^{13}R^{12} \\ R^{12}R^{23} = R^{23}R^{12} \end{cases} \tag{5}$$

holds in $\mathrm{End}_k(M \otimes M \otimes M)$.

Remarks 2.2 1. If R is a solution of the Long equation, then R satisfies the equation

$$[R^{12}, R^{13} + R^{23}] = 0.$$

Hence, any symmetric solution R (i.e. $R^{12} = R^{21}$) of the Long equation is a solution for the integrability condition of the Knizhnik-Zamolodchikov equation (see [4] or [10]).

2. Let M be a finite dimensional vector space and $\{m_1, \cdots, m_n\}$ a basis of M. Let $R \in \mathrm{End}_k(M \otimes M)$ given by

$$R(m_v \otimes m_u) = \sum_{i,j} x_{uv}^{ji} m_i \otimes m_j,$$

for all u, $v = 1, \cdots, n$, where $(x_{uv}^{ji})_{i,j,u,v}$ is a family of scalars of k. Then R is a solution of the Long equation if and only if the following two equations hold:

$$\sum_v x_{kv}^{ji} x_{ql}^{pv} = \sum_\alpha x_{kl}^{j\alpha} x_{q\alpha}^{pi} \tag{6}$$

$$\sum_v x_{kv}^{ji} x_{lq}^{vp} = \sum_\alpha x_{kl}^{j\alpha} x_{\alpha q}^{ip} \tag{7}$$

for all i, j, k, l, p, $q = 1, \cdots, n$.

3. The second equation of our system, namely $R^{12}R^{23} = R^{23}R^{12}$ (called the $\mathcal{D}$-equation), which is obtained from the quantum Yang-Baxter equation $R^{12}R^{13}R^{23} = R^{23}R^{13}R^{12}$ by deleting the middle term from both sides, was studied in [8]. For a bialgebra H, let ${}_H\mathcal{L}^H$ be the category of H-dimodules. In [8] we proved that in the finite dimensional case the maps

$$R_{(M,\cdot,\rho)} : M \otimes M \to M \otimes M, \quad R_{(M,\cdot,\rho)}(m \otimes n) := \sum n_{<1>} \cdot m \otimes n_{<0>},$$

where $(M, \cdot, \rho)$ is a $D(R)$-dimodule for some bialgebra $D(R)$, describle all the solutions of the $\mathcal{D}$-equation.

Examples 2.3 1. Suppose that $R \in \mathrm{End}_k(M \otimes M)$ is bijective. Then, R is a solution of the Long equation if and only if R^{-1} is.

2. Let $(m_i)_{i \in I}$ be a basis of M and $(a_{ij})_{i,j \in I}$ be a family of scalars of k. Then, $R : M \otimes M \to M \otimes M$, $R(m_i \otimes m_j) = a_{ij} m_i \otimes m_j$, for all i, $j \in I$, is a solution of the Long equation. In particular, the identity map $Id_{M \otimes M}$ is a solution of the Long equation.

3. Let M be a finite dimensional vector space and u an automorphism of M. If R is a solution of the Long equation, then ${}^uR := (u \otimes u)R(u \otimes u)^{-1}$ is also a solution of the Long equation.

4. Let A be a k-algebra, $(a_i)_{i=1,\cdots,n}$ be a family of elements of A and $(M, \cdot)$ a left A-module. Then

$$R : M \otimes M \to M \otimes M, \quad R(l \otimes m) := \sum_{i=1}^n l \otimes a_i \cdot m$$

for all l, $m \in M$, is a solution of the Long equation.

5. Let A be a k-algebra and $R = \sum R^1 \otimes R^2 \in A \otimes A$ such that

$$\sum R^1 a \otimes R^2 = \sum aR^1 \otimes R^2 \tag{8}$$

for all $a \in A$. If $(M, \cdot)$ is a left A-module, then the homothety

$$\mathcal{R} : M \otimes M \to M \otimes M, \quad \mathcal{R}(m \otimes n) = R \cdot (m \otimes n) = \sum R^1 \cdot m \otimes R^2 \cdot n,$$

is a solution of the Long equation.

6. Let f, $g \in \operatorname{End}_k(M)$ such that $fg = gf$. Then, $R := f \otimes g$ is a solution of the Long equation.

In particular, let M be a two-dimensional vector space with $\{m_1, m_2\}$ a basis. Let f, $g \in \operatorname{End}_k(M)$ such that, with respect to the given basis, they are:

$$f = \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}, \quad g = \begin{pmatrix} b & c \\ 0 & b \end{pmatrix}$$

where a, b, c are scalars of k. Then, $R = f \otimes g$, with respect to the ordered basis $\{m_1 \otimes m_1, m_1 \otimes m_2, m_2 \otimes m_1, m_2 \otimes m_2\}$ of $M \otimes M$, is given by

$$R = \begin{pmatrix} ab & ac & b & c \\ 0 & ab & 0 & b \\ 0 & 0 & ab & ac \\ 0 & 0 & 0 & ab \end{pmatrix} \tag{9}$$

and R is a solution of the Long equation.

7. Let G be an abelian group and $(M, \cdot)$ be a $k[G]$-module. Suppose that there exists $\{M_\sigma \mid \sigma \in G\}$ a family of $k[G]$-submodules of M such that $M = \oplus_{\sigma \in G} M_\sigma$. If $m \in M$, then m is a finite sum of homogenous elements $m = \sum m_\sigma$. The map

$$R : M \otimes M \to M \otimes M, \quad R(n \otimes m) = \sum_\sigma \sigma \cdot n \otimes m_\sigma, \ \forall n, m \in M \tag{10}$$

is a solution of the Long equation.

In [3], for an n-dimensional vector space M, a certain operator $R_\phi : M \otimes M \to M \otimes M$ is associated to any function $\phi : \{1, \cdots, n\} \to \{1, \cdots, n\}$, with $\phi^2 = \phi$. This operator R_ϕ is a solution for what we called in [6] the Hopf equation

$$R^{23} R^{13} R^{12} = R^{12} R^{23}$$

We shall modify the operator R_ϕ in order to make it a solution for the Long equation.

Proposition 2.4 *Let M be an n-dimensional vector space with $\{m_1, \cdots, m_n\}$ a basis and $\phi : \{1, \cdots, n\} \to \{1, \cdots, n\}$ a function with $\phi^2 = \phi$. Let*

$$R^\phi : M \otimes M \to M \otimes M, \quad R^\phi(m_i \otimes m_j) = \delta_{ij} \delta_{i \in Im(\phi)} \sum_{a,b \in \phi^{-1}(i)} m_a \otimes m_b$$

for all $i,\ j = 1, \cdots, n$*. Then* $R := R^{\phi}$ *is a symmetric solution for the equation*

$$R^{12}R^{13} = R^{13}R^{12} = R^{12}R^{23} = R^{23}R^{12}.$$

In particular, R^{ϕ} *is a solution of the Long equation.*

Proof For $i,\ j,\ k = 1, \cdots, n$ we have:

$$\begin{aligned}
R^{12}R^{13}(m_i \otimes m_j \otimes m_k) &= R^{12}\Big(\delta_{ik}\delta_{i\in Im(\phi)} \sum_{a,b\in\phi^{-1}(i)} m_a \otimes m_j \otimes m_b\Big) \\
&= \delta_{ik}\delta_{i\in Im(\phi)} \sum_{a,b\in\phi^{-1}(i)} \delta_{aj}\delta_{a\in Im(\phi)} \sum_{c,d\in\phi^{-1}(a)} m_c \otimes m_d \otimes m_b \\
&= \delta_{ik}\delta_{i\in Im(\phi)}\delta_{j\in\phi^{-1}(i)}\delta_{j\in Im(\phi)} \sum_{b\in\phi^{-1}(i),c,d\in\phi^{-1}(j)} m_c \otimes m_d \otimes m_b \\
\text{(using } \phi^2 = \phi \text{)} &= \delta_{ik}\delta_{ij}\delta_{i\in\phi^{-1}(i)} \sum_{b,c,d\in\phi^{-1}(i)} m_c \otimes m_d \otimes m_b
\end{aligned}$$

and

$$\begin{aligned}
R^{13}R^{12}(m_i \otimes m_j \otimes m_k) &= R^{13}\Big(\delta_{ij}\delta_{i\in Im(\phi)} \sum_{a,d\in\phi^{-1}(i)} m_a \otimes m_d \otimes m_k\Big) \\
&= \delta_{ij}\delta_{i\in Im(\phi)} \sum_{a,d\in\phi^{-1}(i)} \delta_{ak}\delta_{a\in Im(\phi)} \sum_{c,b\in\phi^{-1}(a)} m_c \otimes m_d \otimes m_b \\
&= \delta_{ij}\delta_{i\in Im(\phi)}\delta_{k\in\phi^{-1}(i)}\delta_{k\in Im(\phi)} \sum_{d\in\phi^{-1}(i),b,c\in\phi^{-1}(k)} m_c \otimes m_d \otimes m_b \\
\text{(using } \phi^2 = \phi \text{)} &= \delta_{ij}\delta_{ik}\delta_{i\in\phi^{-1}(i)} \sum_{b,c,d\in\phi^{-1}(i)} m_c \otimes m_d \otimes m_b
\end{aligned}$$

i.e. $R^{12}R^{13} = R^{13}R^{12}$. The proof of the second identity ($R^{12}R^{23} = R^{23}R^{12}$) is left to the reader.

On the other hand the family of scalars (x_{uv}^{ji}) which define R^{ϕ} are given by $x_{uv}^{ji} = \delta_{uv}\delta_{\phi(i)v}\delta_{\phi(j)v}$ for all $i,\ j,\ u,\ v = 1, \cdots, n$. Hence, $x_{uv}^{ji} = x_{vu}^{ji}$, i.e. R is symmetric. $\square$

3 Long bialgebras

Recall from [8] the following definition

Definition 3.1 *Let* C *be a coalgebra. A* k*-bilinear map* $\sigma : C \otimes C \to k$ *is called a strong* $\mathcal{D}$*-map if*

$$\sum \sigma(c_{(1)} \otimes d)c_{(2)} = \sum \sigma(c_{(2)} \otimes d)c_{(1)} \tag{11}$$

for all $c,\ d \in C$*.*

Proposition 3.2 *Let C be a coalgebra and $\sigma : C \otimes C \to k$ a strong $\mathcal{D}$-map. Let (M, ρ) be a right C-comodule. Then, the special map*

$$R_\sigma : M \otimes M \to M \otimes M, \quad R_\sigma(m \otimes n) = \sum \sigma(m_{<1>} \otimes n_{<1>}) m_{<0>} \otimes n_{<0>}$$

is a solution of the Long equation.

Proof Let $R = R_\sigma$. Then, the fact that $R^{12}R^{23} = R^{23}R^{12}$ is Proposition 4.3 of [8]. We show that $R^{12}R^{13} = R^{13}R^{12}$. For l, m, $n \in M$ we have:

$$\begin{aligned} R^{12}R^{13}(l \otimes m \otimes n) &= R^{12}\Big(\sum \sigma(l_{<1>} \otimes n_{<1>}) l_{<0>} \otimes m \otimes n_{<0>}\Big) \\ &= \sum \sigma(l_{<2>} \otimes n_{<1>}) \sigma(l_{<1>} \otimes m_{<1>}) l_{<0>} \otimes m_{<0>} \otimes n_{<0>} \end{aligned}$$

and

$$\begin{aligned} R^{13}R^{12}(l \otimes m \otimes n) &= R^{13}\Big(\sum \sigma(l_{<1>} \otimes m_{<1>}) l_{<0>} \otimes m_{<0>} \otimes n\Big) \\ &= \sum \sigma(l_{<2>} \otimes m_{<1>}) \sigma(l_{<1>} \otimes n_{<1>}) l_{<0>} \otimes m_{<0>} \otimes n_{<0>} \\ &= \sum \sigma\Big(\underline{\sigma(l_{<1>(1)} \otimes n_{<1>}) l_{<1>(2)}} \otimes m_{<1>}\Big) l_{<0>} \otimes m_{<0>} \otimes n_{<0>} \\ \text{(using (11))} \quad &= \sum \sigma\Big(\underline{\sigma(l_{<1>(2)} \otimes n_{<1>}) l_{<1>(1)}} \otimes m_{<1>}\Big) l_{<0>} \otimes m_{<0>} \otimes n_{<0>} \\ &= \sum \sigma(l_{<2>} \otimes n_{<1>}) \sigma(l_{<1>} \otimes m_{<1>}) l_{<0>} \otimes m_{<0>} \otimes n_{<0>} \end{aligned}$$

i.e. $R^{12}R^{13} = R^{13}R^{12}$. Hence, R_σ is a solution of the Long equation. □

Now, we shall introduce a new class of bialgebras which play for the Long equation the same role as the co-quasitriangular (or braided) bialgebras do for the quantum Yang-Baxter equation.

Definition 3.3 *A Long bialgebra is a pair (H, σ), where H is a bialgebra and $\sigma : H \otimes H \to k$ is a k-linear map such that the following conditions are fulfilled:*

$(L1)$ $\quad \sum \sigma(x_{(1)} \otimes y) x_{(2)} = \sum \sigma(x_{(2)} \otimes y) x_{(1)}$

$(L2)$ $\quad \sigma(x \otimes 1) = \varepsilon(x)$

$(L3)$ $\quad \sigma(x \otimes yz) = \sum \sigma(x_{(1)} \otimes y) \sigma(x_{(2)} \otimes z)$

$(L4)$ $\quad \sigma(1 \otimes x) = \varepsilon(x)$

$(L5)$ $\quad \sigma(xy \otimes z) = \sum \sigma(y \otimes z_{(1)}) \sigma(x \otimes z_{(2)})$

for all x, y, $z \in H$.

Remark 3.4 We recall that (see, for instance, [4]) a pair (H, σ) is a co-quasitriangular (or braided) bialgebra if σ satisfies the conditions $(B1)$-$(B5)$. The conditions $(B2)$-$(B5)$ are identical with the conditions $(L2)$-$(L5)$. The first condition is

$(B1) \quad \sum \sigma(x_{(1)} \otimes y_{(1)}) y_{(2)} x_{(2)} = \sum \sigma(x_{(2)} \otimes y_{(2)}) x_{(1)} y_{(1)}$

for all $x, y \in H$. If (H, σ) is a co-quasitriangular bialgebra and (M, ρ) is a right H-comodule, then the special map

$$R_\sigma : M \otimes M \to M \otimes M, \quad R_\sigma(m \otimes n) = \sum \sigma(m_{<1>} \otimes n_{<1>}) m_{<0>} \otimes n_{<0>}$$

is a solution for the quantum Yang-Baxter equation $R^{12}R^{13}R^{23} = R^{23}R^{13}R^{12}$. Conversely, if M is a finite dimensional vector space and R is a solution of the quantum Yang-Baxter equation, then there exists a bialgebra $A(R)$ and a unique k-bilinear map $\sigma : A(R) \otimes A(R) \to k$ such that $(A(R), \sigma)$ is co-quasitriangular, $M \in \mathcal{M}^{A(R)}$ and $R = R_\sigma$ (see [2] or [4]).

We shall now present a few examples of Long bialgebras. More examples will be given after the main result.

Examples 3.5 1. If H is cocommutative, then $(L1)$ holds for any k-linear map $\sigma : H \otimes H \to k$. In particular, if G is a group and $\sigma : G \times G \to k$ is a bicharacter on G, then $(k[G], \sigma)$ is a Long bialgebra.

2. Let $H = k < x, y >$ be the free algebra generated by x and y, with the bialgebra structure given by

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes 1 + x \otimes y, \quad \varepsilon(x) = 1, \quad \varepsilon(y) = 0.$$

Let $\sigma : H \otimes H \to k$ be a k-bilinear map. Then, it is easy to see that (H, σ) is a Long bialgebra if and only if

$$\begin{array}{lll} \sigma(x \otimes 1) = 1, & \sigma(x \otimes x) = 1, & \sigma(x \otimes y) = 0, \\ \sigma(y \otimes 1) = 0, & \sigma(y \otimes x) = 0, & \sigma(y \otimes y) = 0. \end{array}$$

3. Let $H = k < x, y, z >$ be the free algebra generated by x, y, z, with the bialgebra structure given by

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y, \quad \Delta(z) = x \otimes z + z \otimes y$$

$$\varepsilon(x) = \varepsilon(y) = 1, \quad \varepsilon(z) = 0.$$

Let $\sigma : H \otimes H \to k$ be a k-bilinear map. Then (H, σ) is a Long bialgebra if and only if there exist a, b, c scalars of k such that

$$\sigma(x \otimes x) = \sigma(y \otimes x) = a, \quad \sigma(x \otimes y) = \sigma(y \otimes y) = b, \quad \sigma(x \otimes z) = \sigma(y \otimes z) = c,$$

$$\sigma(z \otimes x) = \sigma(z \otimes y) = \sigma(z \otimes z) = 0.$$

4. Let H_4 be Sweedler's 4-dimensional Hopf algebra, i.e.

$$H_4 = k < x, y, z \mid x^2 = 1, \ y^2 = 0, \ xy = z, \ xz = -zx = y >$$

with

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes x + 1 \otimes y, \quad \Delta(z) = x \otimes z + z \otimes 1$$

$$\varepsilon(x) = 1, \quad \varepsilon(y) = \varepsilon(z) = 0.$$

It is well known that H_4 is a co-quasitriangular bialgebra (see, for example, [1]). We shall prove that there exists no $\sigma : H_4 \otimes H_4 \to k$ such that (H_4, σ) is a Long bialgebra.

Suppose that there exists $\sigma : H_4 \otimes H_4 \to k$ such that (H_4, σ) is a Long bialgebra. It follows from $(L1)$ that

$$\sigma(y \otimes h)x + \sigma(1 \otimes h)y = \sigma(x \otimes h)y + \sigma(y \otimes h)1_H$$

for all $h \in H$. As $\{1, x, y, z\}$ is a basis of H_4, we get that $\sigma(x \otimes h) = \sigma(1 \otimes h)$ and $\sigma(y \otimes h) = 0$ for all $h \in H$. In particular, $\sigma(x \otimes x) = 1$ and $\sigma(x \otimes z) = \sigma(x \otimes y) = 0$. It follows that

$$0 = \sigma(x \otimes y) = \sigma(x \otimes xz) = \sigma(x \otimes x) + \sigma(x \otimes z)$$

hence $\sigma(x \otimes z) = -1$, contradiction.

Proposition 3.6 *Let H be a bialgebra and $\sigma : H \otimes H \to k$ a k-bilinear map which satisfies $(L3)$ and $(L5)$. Suppose that $(L1)$ holds for a system of generators of H as an algebra. Then $(L1)$ holds for all elements of H.*

Proof Let x, y, $z \in H$ be three elements among the generators of H. It is enough to prove that $(L1)$ holds for (x, yz) and (xy, z). We have:

$$\begin{array}{rcl}
\sum \sigma(x_{(1)} \otimes yz)x_{(2)} & = & \sum \sigma(x_{(1)(1)} \otimes y)\sigma(x_{(1)(2)} \otimes z)x_{(2)} \\
 & = & \sum \sigma(x_{(1)} \otimes y)\underline{\sigma(x_{(2)} \otimes z)x_{(3)}} \\
\text{(using (L1))} & = & \sum \sigma(x_{(1)} \otimes y)\sigma(x_{(3)} \otimes z)x_{(2)} \\
 & = & \sum \underline{\sigma(x_{(1)} \otimes y)x_{(2)}}\sigma(x_{(3)} \otimes z) \\
\text{(using (L1))} & = & \sum \sigma(x_{(2)} \otimes y)\sigma(x_{(3)} \otimes z)x_{(1)} \\
\text{(using (L3))} & = & \sum \sigma(x_{(2)} \otimes yz)x_{(1)}
\end{array}$$

i.e. $(L1)$ holds for (x, yz). On the other hand

$$\begin{array}{rcl}
\sum \sigma((xy)_{(1)} \otimes z)(xy)_{(2)} & = & \sum \sigma(x_{(1)}y_{(1)} \otimes z)x_{(2)}y_{(2)} \\
 & = & \sum \sigma(y_{(1)} \otimes z_{(1)})\underline{\sigma(x_{(1)} \otimes z_{(2)})x_{(2)}}y_{(2)} \\
\text{(using (L1))} & = & \sum \sigma(x_{(2)} \otimes z_{(2)})x_{(1)}\underline{\sigma(y_{(1)} \otimes z_{(1)})y_{(2)}} \\
\text{(using (L1))} & = & \sum \sigma(x_{(2)} \otimes z_{(2)})x_{(1)}\sigma(y_{(2)} \otimes z_{(1)})y_{(1)} \\
\text{(using (L5))} & = & \sum \sigma(x_{(2)}y_{(2)} \otimes z)x_{(1)}y_{(1)} \\
 & = & \sum \sigma((xy)_{(2)} \otimes z)(xy)_{(1)}
\end{array}$$

and the proof is complete now. □

We recall that if M is a right H-comodule over a co-quasitriangular bialgebra (H, σ), then M can be viwed as an object of ${}_H\mathcal{YD}^H$, the category of Yetter-Drinfel'd modules, where the structure of left H-module on M is induced by σ:

$$h \cdot m = \sum \sigma(m_{<1>} \otimes h) m_{<0>}$$

for all $h \in H$, $m \in M$. In the next proposition we shall prove that any right H-comodule M over a Long bialgebra (H, σ) can be viewed as an H-dimodule.

Proposition 3.7 *Let (H, σ) be a Long bialgebra and (M, ρ) be a right H-comodule. Then, the left action of H on M given by*

$$h \cdot m = \sum \sigma(m_{<1>} \otimes h) m_{<0>}$$

for all $h \in H$, $m \in M$, makes $(M, \cdot, \rho)$ an H-dimodule.

Proof The fact that $(M, \cdot)$ is a left H-module follows from $(L2)$ and $(L3)$. We shall prove that the compatibility condition (4) holds. For $h \in H$ and $m \in M$ we have

$$\begin{aligned}
\rho(h \cdot m) &= \rho\Big(\sum \sigma(m_{<1>} \otimes h) m_{<0>}\Big) \\
&= \sum \sigma(m_{<1>} \otimes h) m_{<0><0>} \otimes m_{<0><1>} \\
&= \sum m_{<0>} \otimes \sigma(m_{<1>(2)} \otimes h) m_{<1>(1)} \\
\text{(using (L1))} \quad &= \sum m_{<0>} \otimes \sigma(m_{<1>(1)} \otimes h) m_{<1>(2)} \\
&= \sum \sigma(m_{<1>} \otimes h) m_{<0>} \otimes m_{<2>} \\
&= \sum h \cdot m_{<0>} \otimes m_{<1>}
\end{aligned}$$

i.e. $(M, \cdot, \rho)$ is an H-dimodule. □

From Proposition 3.2 we obtain that if (H, σ) is a Long bialgebra and M is a right H-comodule, then the natural map R_σ is a solution of the Long equation. Now, we shall prove the main result of this paper: in the finite dimensional case any solution of the Long equation arises in this way.

Theorem 3.8 *Let M be a finite dimensional vector space and $R \in \mathrm{End}_k(M \otimes M)$ be a solution for the Long equation. Then there exists a bialgebra $L(R)$ and a unique k-bilinear map*

$$\sigma : L(R) \otimes L(R) \to k$$

such that $(L(R), \sigma)$ is a Long bialgebra, M is a right $L(R)$-comodule and $R = R_\sigma$.

Furthermore, if R is bijective, then σ is invertible in convolution.

Proof Suppose that $\dim_k(M) = n$ and let $\{m_1, \cdots, m_n\}$ be a basis for M. Let $(x_{uv}^{ji})_{i,j,u,v}$ be a family of scalars of k such that

$$R(m_v \otimes m_u) = \sum_{i,j} x_{uv}^{ji} m_i \otimes m_j \tag{12}$$

for all u, $v = 1, \cdots, n$.

Let $\mathcal{M}^n(k)$, be the comatrix coalgebra of order n with $\{c_{ij} \mid i, j = 1, \cdots, n\}$ a k-basis and $(T(\mathcal{M}^n(k)), \Delta, \varepsilon, \mu, 1)$ the unique bialgebra structure on the tensor algebra $(T(\mathcal{M}^n(k)), \mu, 1)$ whose comultiplication Δ and counit ε extend the comultiplication and the counit from $\mathcal{M}^n(k)$. Then M has a unique right $T(\mathcal{M}^n(k))$-comodule structure $\rho : M \to M \otimes T(\mathcal{M}^n(k))$ such that

$$\rho(m_l) = \sum_{v=1}^{n} m_v \otimes c_{vl} \tag{13}$$

for all $l = 1, \cdots, n$. Let $o(i, j, k, l)$ be the obstruction

$$o(i, j, k, l) := \sum_{v} x_{kv}^{ji} c_{vl} - \sum_{\alpha} x_{kl}^{j\alpha} c_{i\alpha} \tag{14}$$

for all i, j, k, $l = 1, \cdots, n$. Let $I(R)$ be the two-sided ideal of $T(\mathcal{M}^n(k))$ generated by all $o(i, j, k, l)$. Then, $I(R)$ is also a coideal of $T(\mathcal{M}^n(k))$, i.e. it is a biideal. This follows from the formulas

$$\Delta(o(i, j, k, l)) = \sum_{u} \Big(o(i, j, k, u) \otimes c_{ul} + c_{iu} \otimes o(u, j, k, l) \Big)$$

for all i, j, k, $l = 1, \cdots, n$ (see equation (16) of [8]). We now define

$$L(R) = T(\mathcal{M}^n(k))/I(R)$$

which is a bialgebra. M has a right $L(R)$-comodule structure via the natural projection $T(\mathcal{M}^n(k)) \to L(R)$, i.e.

$$\rho(m_l) = \sum_{v=1}^{n} m_v \otimes \overline{c_{vl}}$$

for all $l = 1, \cdots, n$.

First we shall prove the uniqueness of σ. Let $\sigma : L(R) \otimes L(R) \to k$ such that $(L(R), \sigma)$ is a Long bialgebra and $R = R_\sigma$. Let u, $v = 1, \cdots, n$. Then

$$R_\sigma(m_v \otimes m_u) == \sum_{i,j} \sigma(\overline{c_{iv}} \otimes \overline{c_{ju}}) m_i \otimes m_j$$

Hence $R_\sigma(m_v \otimes m_u) = R(m_v \otimes m_u)$ gives us

$$\sigma(\overline{c_{iv}} \otimes \overline{c_{ju}}) = x_{uv}^{ji} \tag{15}$$

for all i, j, u, $v = 1, \cdots, n$. As $L(R)$ is generated as an algebra by $(\overline{c_{ij}})$, the relations (15) with $(L2) - (L5)$ ensure the uniqueness of σ.

Now we shall prove the existence of σ. First we define

$$\sigma_0 : \mathcal{M}^n(k) \otimes \mathcal{M}^n(k) \to k, \qquad \sigma_0(c_{iv} \otimes c_{ju}) = x^{ji}_{uv}$$

for all i, j, u, $v = 1, \cdots, n$. Then, we extend σ_0 to a map

$$\sigma_1 : T(\mathcal{M}^n(k)) \otimes T(\mathcal{M}^n(k)) \to k$$

such that $(L2) - (L5)$ hold. We shall prove that σ_1 factorizes to a map

$$\sigma : L(R) \otimes L(R) \to k$$

i.e.

$$\sigma_1\Big(T(\mathcal{M}^n(k)) \otimes o(i,j,k,l)\Big) = \sigma_1\Big(o(i,j,k,l) \otimes T(\mathcal{M}^n(k))\Big) = 0.$$

For i, j, k, l, p, $q = 1, \cdots, n$ we have:

$$\begin{aligned} \sigma_1(c_{pq} \otimes o(i,j,k,l)) &= \sum_v x^{ji}_{kv}\sigma_1(c_{pq} \otimes c_{vl}) - \sum_\alpha x^{j\alpha}_{kl}\sigma_1(c_{pq} \otimes c_{i\alpha}) \\ &= \sum_v x^{ji}_{kv} x^{vp}_{lq} - \sum_\alpha x^{j\alpha}_{kl} x^{ip}_{\alpha q} \\ \text{(from (7))} &= 0 \end{aligned}$$

and

$$\begin{aligned} \sigma_1(o(i,j,k,l) \otimes c_{pq}) &= \sum_v x^{ji}_{kv}\sigma_1(c_{vl} \otimes c_{pq}) - \sum_\alpha x^{j\alpha}_{kl}\sigma_1(c_{i\alpha} \otimes c_{pq}) \\ &= \sum_v x^{ji}_{kv} x^{pv}_{ql} - \sum_\alpha x^{j\alpha}_{kl} x^{pi}_{q\alpha} \\ \text{(from (6))} &= 0 \end{aligned}$$

Hence, we have constructed $\sigma : L(R) \otimes L(R) \to k$ such that $(L2) - (L5)$ hold and $R = R_\sigma$. It remains to prove that σ satisfies $(L1)$. Using proposition 3.6, it is enough to show that $(L1)$ holds on the generators. Let $x = \overline{c_{ij}}$, $y = \overline{c_{pq}}$. We have:

$$\sum \sigma(x_{(1)} \otimes y)x_{(2)} = \sum_v \sigma(\overline{c_{iv}} \otimes \overline{c_{pq}})\, \overline{c_{vj}} = \sum_v x^{pi}_{qv}\overline{c_{vj}}$$

and

$$\sum \sigma(x_{(2)} \otimes y)x_{(1)} = \sum_\alpha \sigma(\overline{c_{\alpha j}} \otimes \overline{c_{pq}})\, \overline{c_{i\alpha}} = \sum_\alpha x^{p\alpha}_{qj}\overline{c_{i\alpha}}.$$

Hence,

$$\sum \sigma(x_{(1)} \otimes y)\, x_{(2)} - \sum \sigma(x_{(2)} \otimes y)\, x_{(1)} = \overline{o(i,p,q,j)} = 0,$$

i.e. $(L1)$ holds. Suppose now that R is bijective and let $S = R^{-1}$. Let (y^{ji}_{uv}) be a family of scalars of k such that

$$S(m_v \otimes m_u) = \sum_{i,j} y^{ji}_{uv} m_i \otimes m_j,$$

for all u, $v = 1, \cdots, n$. We define

$$\sigma' : L(R) \otimes L(R) \to k, \quad \sigma'(\overline{c_{iv}} \otimes \overline{c_{ju}}) := y^{ji}_{uv}$$

for all $i,\ j,\ u,\ v = 1, \cdots, n$. Now it is routine to see that σ' is well defined and it is the inverse of σ. $\square$

Remark 3.9 Let M be a finite dimensional vector space and $R \in \mathrm{End}_k(M \otimes M)$ be a solution for the Long equation. Using theorem 3.8 and proposition 3.7, we obtain that M has a structure of a $L(R)$-dimodule $(M, \cdot, \rho)$ and $R = R_{(M,\cdot,\rho)}$.

Examples 3.10 1. Let $a,\ b,\ c \in k$ such that $(b, c) \neq 0$ and $R \in \mathcal{M}_4(k)$ given by the formula (9). Then, R is a solution of the Long equation. The bialgebra $L(R)$ can be described as follows:

- as an algebra $L(R) = k < x, y >$, the free algebra generated by x and y.
- the comultiplication Δ and the counity ε are given by

$$\Delta(x) = x \otimes x, \quad \Delta(y) = x \otimes y + y \otimes x, \quad \varepsilon(x) = 1, \quad \varepsilon(y) = 0.$$

Indeed, if we write

$$R(m_v \otimes m_u) = \sum_{i,j=1}^{2} x_{uv}^{ji} m_i \otimes m_j$$

we get that, among the elements (x_{uv}^{ji}), the only nonzero elements are:

$$x_{11}^{11} = ab, \quad x_{21}^{11} = ac, \quad x_{21}^{21} = ab, \quad x_{12}^{11} = b,$$

$$x_{12}^{12} = ab, \quad x_{21}^{11} = c, \quad x_{22}^{21} = b, \quad x_{22}^{12} = ac, \quad x_{22}^{22} = ab$$

The sixteen relations $o(i, j, k, l) = 0$, written in lexicographic order according to (i, j, k, l), starting with $(1, 1, 1, 1)$, are

$$abc_{11} + bc_{21} = abc_{11}, \quad abc_{12} + bc_{22} = bc_{11} + abc_{12}$$

$$acc_{11} + cc_{21} = acc_{11}, \quad acc_{12} + cc_{22} = cc_{11} + acc_{12}$$

$$0 = 0, \quad 0 = 0, \quad abc_{11} + bc_{21} = abc_{11}, \quad abc_{12} + bc_{22} = bc_{11} + abc_{12}$$

$$abc_{21} = abc_{21}, \quad abc_{22} = abc_{22}, \quad acc_{21} = acc_{21}, \quad acc_{22} = cc_{21} + acc_{22}$$

$$0 = 0, \quad 0 = 0, \quad abc_{21} = abc_{21}, \quad abc_{22} = abc_{22} + bc_{21}$$

Only the four relations below remain:

$$bc_{21} = 0, \quad bc_{22} = bc_{11}, \quad cc_{21} = 0, \quad cc_{22} = cc_{11}$$

As $(b, c) \neq (0, 0)$, there are only two linear independent relations:

$$c_{21} = 0, \qquad c_{22} = c_{11}$$

Now, if we denote $c_{11} = x$, $c_{12} = y$, the above description of $L(R)$ follows. Moreover, $L(R)$ can be viewed as a Long bialgebra, where $\sigma : k < x, y > \otimes k < x, y > \to k$ is given by:

$$\begin{aligned} &\sigma(x \otimes 1) = 1, \quad \sigma(x \otimes x) = ab, \quad \sigma(x \otimes y) = c, \\ &\sigma(y \otimes 1) = 0, \quad \sigma(y \otimes x) = b, \quad \sigma(y \otimes y) = 0. \end{aligned}$$

2. Let a, $b \in k$ and $R \in \mathcal{M}_4(k)$ given by

$$R = \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Then R is a solution of the Long equation, as $R = f \otimes g$, where f and g are given by

$$f = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \qquad g = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$$

i.e. $fg = gf$. We shall describe the bialgebra $L(R)$. Suppose $(a, b) \neq 0$ (otherwise $R = 0$ and $L(R) = T(\mathcal{M}^4(k))$). Among the sixteen relations $o(i, j, k, l) = 0$, the only linear independent ones are:

$$ac_{12} = ac_{21} = bc_{12} = bc_{21} = 0$$

As $(a, b) \neq 0$, we obtain $c_{12} = c_{21} = 0$. Now, if we denote $c_{11} = x$, $c_{22} = y$, we get that the bialgebra $L(R)$ has the following description:

- as an algebra, $L(R) = k < x, y >$, the free algebra generated by x and y.
- the comultiplication Δ and the counity ε are given in such a way that x and y are groupal elements

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y \quad \varepsilon(x) = \varepsilon(y) = 1.$$

Moreover, $k < x, y >$ is a Long bialgebra as follows:

$$\begin{array}{lll} \sigma(x \otimes 1) = 1, & \sigma(x \otimes x) = a, & \sigma(x \otimes y) = b, \\ \sigma(y \otimes 1) = 1, & \sigma(y \otimes x) = 0 & \sigma(y \otimes y) = 0. \end{array}$$

3. Let M be an n-dimensional vector space with $\{m_1, \cdots, m_n\}$ a basis and $\phi : \{1, \cdots, n\} \to \{1, \cdots, n\}$ a function with $\phi^2 = \phi$. We have proven that the map

$$R^\phi : M \otimes M \to M \otimes M, \quad R^\phi(m_i \otimes m_j) = \delta_{ij} \delta_{i \in Im(\phi)} \sum_{a,b \in \phi^{-1}(i)} m_a \otimes m_b \tag{16}$$

for all i, $j = 1, \cdots, n$, is a solution of the Long equation. Below, we shall describe the bialgebra $L(R^\phi)$. From equation (16), the scalars (x_{uv}^{ji}) which define R^ϕ are given by

$$x_{uv}^{ji} = \delta_{uv} \delta_{\phi(i)v} \delta_{\phi(j)v}$$

for all i, j, u, $v = 1, \cdots, n$. The relations $o(i, j, k, l) = 0$ are

$$\sum_v \delta_{kv} \delta_{\phi(i)v} \delta_{\phi(j)v} c_{vl} = \delta_{kl} \delta_{\phi(j)l} \sum_\alpha \delta_{\phi(\alpha)l} c_{i\alpha}$$

or equivalent to

$$\delta_{\phi(i)k} \delta_{\phi(j)k} c_{kl} = \delta_{kl} \delta_{\phi(j)l} \sum_\alpha \delta_{\phi(\alpha)l} c_{i\alpha}$$

for all $i,\ j,\ u,\ v = 1, \cdots, n$. All the relations $o(i,j,k,l) = 0$ are identities $0 = 0$, with the exception of the following three types of relations:

$$\begin{cases} o(i,j,\phi(j),l) = 0, & \text{for all} \quad l \neq \phi(j),\ \phi(i) = \phi(j) \\ o(i,j,\phi(j),\phi(j)) = 0, & \text{for all} \quad \phi(i) \neq \phi(j) \\ o(i,j,\phi(j),\phi(j)) = 0, & \text{for all} \quad \phi(i) = \phi(j) \end{cases} \tag{17}$$

which give us the following description of the bialgebra $L(R^\phi)$: as an algebra, $L(R^\phi)$ is the free algebra generated by (c_{ij}) with the relations:

$$\begin{cases} c_{\phi(j)l} = 0, & \text{for all} \quad l \neq \phi(j) \\ \sum_{\alpha \in \phi^{-1}(\phi(j))} c_{i\alpha} = 0, & \text{for all} \quad \phi(i) \neq \phi(j) \\ \sum_{\alpha \in \phi^{-1}(\phi(i))} c_{i\alpha} = c_{\phi(i)\phi(i)}, & \text{for all} \quad i = 1, \cdots, n \end{cases} \tag{18}$$

The comultiplication and the counity of $L(R^\phi)$ are given in such a way that the matrix (c_{ij}) is comultiplicative. The bialgebra $L(R^\phi)$ is a Long bialgebra with

$$\sigma : L(R^\phi) \otimes L(R^\phi) \to k, \quad \sigma(c_{iv} \otimes c_{ju}) = \delta_{uv}\delta_{\phi(i)v}\delta_{\phi(j)v}$$

for all $i,\ j,\ u,\ v = 1, \cdots, n$.

4. In the next three examples we point out particular cases of the previous example. Let ϕ the identity map, i.e. $\phi(i) = i$ for all $i = 1, \cdots, n$. Then $L(R^\phi) = k < x_1, \cdots, x_n >$, the free algebra generated by $x_1, \cdots, x_n$, and the coalgebra structure is given in such a way that x_i is a group-like element, i.e. $\Delta(x_i) = x_i \otimes x_i$ and $\varepsilon(x_i) = 1$ for all $i = 1, \cdots, n$. Indeed, from the relations (18), only

$$c_{jl} = 0, \forall l \neq j$$

remains. If we denote $c_{ii} = x_i$ the conclusion follows. We note that the bialgebra $k < x_1, \cdots, x_n >$ is a Long bialgebra with $\sigma : k < x_1, \cdots, x_n > \otimes k < x_1, \cdots, x_n > \to k$ given by

$$\sigma(x_i \otimes x_j) = \delta_{ij}$$

for all $i,\ j = 1, \cdots, n$.

5. Let $n = 4$ and ϕ given by

$$\phi(1) = 1, \quad \phi(2) = \phi(3) = \phi(4) = 2.$$

Then the relations (18) are:

$$c_{1l} = c_{2j} = c_{i1} = 0, \quad c_{32} + c_{33} + c_{34} = c_{22}, \quad c_{42} + c_{43} + c_{44} = c_{22}$$

for all $l \neq 1$, $j \neq 2$, $i \neq 1$. Now, if we denote $c_{11} = x_1$, $c_{22} = x_2$, $c_{32} = x_3$, $c_{33} = x_4$, $c_{42} = x_5$, $c_{44} = x_6$ we obtain the description of the corresponding bialgebra $L(R^\phi)$:

- as an algebra $L(R^\phi) = k < x_i \mid i = 1, \cdots, 6 >$, the free algebra generated by six generators.

- the comultiplication Δ and the counity ε are given by

$$\Delta(x_1) = x_1 \otimes x_1, \quad \Delta(x_2) = x_2 \otimes x_2,$$

$$\Delta(x_3) = x_3 \otimes x_2 + x_4 \otimes x_3 + (x_2 - x_3 - x_4) \otimes x_5, \quad \Delta(x_4) = x_4 \otimes x_4 + (x_2 - x_3 - x_4) \otimes (x_2 - x_5 - x_6),$$

$$\Delta(x_5) = x_5 \otimes x_2 + (x_2 - x_5 - x_6) \otimes x_3 + x_6 \otimes x_5, \quad \Delta(x_6) = (x_2 - x_5 - x_6) \otimes (x_2 - x_3 - x_4) + x_6 \otimes x_6$$

$$\varepsilon(x_1) = \varepsilon(x_2) = \varepsilon(x_4) = \varepsilon(x_6) = 1, \quad \varepsilon(x_3) = \varepsilon(x_5) = 0.$$

6. Let $n = 4$ and ϕ given by

$$\phi(1) = \phi(2) = 2, \quad \phi(3) = \phi(4) = 4.$$

Then the relations (18) are:

$$c_{2l} = c_{4j} = 0, \quad c_{11} + c_{12} = c_{22}, \quad c_{33} + c_{34} = c_{44}, \quad c_{13} + c_{14} = c_{31} + c_{32} = 0$$

for all $l \neq 2$, $j \neq 4$. If we denote $c_{11} = x_1$, $c_{12} = x_2$, $c_{13} = x_3$, $c_{31} = x_4$, $c_{33} = x_5$, $c_{34} = x_6$, we obtain the description of the corresponding bialgebra $L(R^\phi)$:

- as an algebra $L(R^\phi) = k < x_i \mid i = 1, \cdots, 6 >$, the free algebra generated by six generators.
- the comultiplication Δ and the counity ε are given by

$$\Delta(x_1) = x_1 \otimes x_1 + x_3 \otimes x_4, \quad \Delta(x_2) = x_1 \otimes x_2 + x_2 \otimes x_1 + x_2 \otimes x_2 - x_3 \otimes x_4,$$

$$\Delta(x_3) = x_1 \otimes x_3 + x_3 \otimes x_5, \quad \Delta(x_4) = x_4 \otimes x_1 + x_5 \otimes x_4,$$

$$\Delta(x_5) = x_4 \otimes x_3 + x_5 \otimes x_5, \quad \Delta(x_6) = -x_4 \otimes x_3 + x_5 \otimes x_6 + x_6 \otimes x_5 + x_6 \otimes x_6$$

$$\varepsilon(x_1) = \varepsilon(x_5) = 1, \quad \varepsilon(x_2) = \varepsilon(x_3) = \varepsilon(x_4) = \varepsilon(x_6) = 0.$$

References

[1] Y. Doi, Braided bialgebras and quadratic bialgebras, Comm. Algebra, 21(1993), 1731-1749.

[2] L. D. Faddeev, N. Yu Reshetikhin, L. A. Takhtadjian, Quantization of Lie groups and Lie algebras, *Algebra i Analiz* **1**(1989), 178-206.

[3] B. Ion, M. Stanciulescu, Several examples of noncommutative and noncocommutative bialgebras arising from the Hopf equation, Revue Roumaine de Math. Pure and Appl., to appear.

[4] C. Kassel, Quantum groups, Springer Verlag, Berlin, 1995.

[5] F.W. Long, The Brauer group of dimodule algebras, J. Algebra, 31(1974), 559-601.

[6] G. Militaru, The Hopf module category and the Hopf equation, *Comm. in Algebra*, to appear.

[7] G. Militaru, New types of bialgebras arising from the Hopf equation, *Comm. in Algebra*, to appear.

[8] G. Militaru, The Long dimodules category and nonlinear equations, preprint 1997.

[9] D. Radford, Solutions to the quantum Yang-Baxter equation and the Drinfel'd double, *J. Algebra* **161** (1993), 20-32.

[10] S. Shnider, S. Sternberg, Quantum groups-from coalgebras to Drinfel'd algebras; a guided tour, International Press Company, Boston, 1993.